

THE AVERAGE GAP DISTRIBUTION FOR GENERALIZED ZECKENDORF DECOMPOSITIONS

OLIVIA BECKWITH, AMANDA BOWER, LOUIS GAUDET, RACHEL INSOFT, SHIYU LI,
STEVEN J. MILLER, AND PHILIP TOSTESON

ABSTRACT. An interesting characterization of the Fibonacci numbers is that, if we write them as $F_1 = 1$, $F_2 = 2$, $F_3 = 3$, $F_4 = 5, \dots$, then every positive integer can be written uniquely as a sum of non-adjacent Fibonacci numbers. This is now known as Zeckendorf's theorem [Ze], and similar decompositions exist for many other sequences $\{G_n = c_1 G_{n-1} + \dots + c_L G_{n-L}\}$ arising from recurrence relations. Much more is known. Using continued fraction approaches, Lekkerkerker [Lek] proved the average number of summands needed for integers in $[G_n, G_{n+1})$ is on the order of $C_{\text{Lek}} n$ for a non-zero constant; this was improved by others to show the number of summands has Gaussian fluctuations about this mean.

Koloğlu, Kopp, Miller and Wang [KKMW, MW1] recently recast the problem combinatorially, reproving these results. We use this new perspective to investigate the distribution of gaps between summands. We explore the average behavior over all $m \in [G_n, G_{n+1})$ for special choices of the c_i 's, and leave the general case as well as the behavior of gap measures associated to individual m 's to the sequel paper [BILMT]. Specifically, we study the case where each $c_i \in \{0, 1\}$ and there is a g such that there are always exactly $g - 1$ zeros between two non-zero c_i 's; note this includes the Fibonacci, Tribonacci and many other important special cases. We prove there are no gaps of length less than g , and the probability of a gap of length $j > g$ decays geometrically, with the decay ratio equal to the largest eigenvalue of the recurrence relation. These methods are combinatorial and apply to related problems; we end with a discussion of similar results for far-difference (i.e., signed) decompositions.

CONTENTS

1. Introduction	2
2. Kangaroo Recurrences	5
2.1. Preliminary Results	5
2.2. Proof of Theorem 1.4	7
2.3. Approximating $\lambda_{g,\ell}$ and Relevant Probability Ratios	8
3. Far-Difference Representations	9
4. Conclusion and Open Questions	10

Date: August 30, 2012.

2010 Mathematics Subject Classification. 11B39, 11B05 (primary) 65Q30, 60B10 (secondary).

Key words and phrases. Fibonacci numbers, linear recurrences, Zeckendorf's Theorem, far-difference representations, geometric decay.

The third and sixth named author was partially supported by NSF grant DMS0970067 and the remaining authors were partially supported by NSF Grant DMS0850577. It is a pleasure to thank our colleagues from the Williams College 2010, 2011 and 2012 SMALL REU program for many helpful conversations, and to Florian Luca for comments on an earlier draft.

1. INTRODUCTION

A beautiful theorem of Zeckendorf [Ze] states that every positive integer can be written uniquely as a sum of non-adjacent Fibonacci numbers if we label them $F_1 = 1$, $F_2 = 2$, $F_3 = 3$, $F_4 = 5$ and so on; we must use this non-standard ordering as otherwise the decomposition would not be unique. The standard proof of this is through a greedy algorithm, though this does not provide any information about finer questions. Using continued fraction techniques, Lekkerkerker [Lek] proved that the average number of summands needed for decompositions of numbers in $[F_n, F_{n+1})$ is $n/(\phi^2 + 1) + O(1)$, where $\phi = (1 + \sqrt{5})/2$ is the golden mean. These results have been extended to other sequences, specifically $\{G_n\}$ being a positive linear recurrence sequence. This means that there exist positive integers c_i (with c_1, c_L non-zero) such that

$$G_n = c_1 G_{n-1} + \cdots + c_L G_{n-L}, \quad (1.1)$$

as well as rules to specify the first L terms of the sequence and a generalization of the non-adjacency constraint to what is a ‘legal’ decomposition (other authors use the phrase G -ary decomposition for a legal decomposition). Informally, a legal decomposition is one where we cannot use the recurrence relation to replace a linear combination of summands with another summand, and the coefficient of each summand is appropriately bounded; see [MW1, MW2]. In addition to generalizations of Zeckendorf’s and Lekkerkerker’s theorems holding (the average number of summands is $C_{\text{Lek}}n + d$ for some $C_{\text{Lek}} > 0$), the fluctuations of the number of summands above the mean converges to Gaussian behavior; see [BCCSW, DG, FGNPT, Ho, Ke, LT, Len, MW1, MW2, Ste1, Ste2].

In this paper we pursue a related question, the distribution of gaps between summands in generalized Zeckendorf decompositions. We adopt the perspective of Koloğlu, Kopp, Miller and Wang [KKMW, MW1, MW2], which we briefly review. They proved Gaussian behavior by recasting the problem combinatorially, writing down explicit formulas for the number of $m \in [G_n, G_{n+1})$ with exactly k summands through the cookie (or stars and bars) problem. Specifically, in the Fibonacci case consider all integers $m \in [F_n, F_{n+1})$ with exactly $k + 1$ summands. If these summands are $F_{i_1}, F_{i_2}, \dots, F_{i_{k+1}=n}$ then the non-adjacency condition implies the gaps are $d_j := i_j - i_{j-1} \geq 2$ for $j \geq 2$ and $d_1 := i_1 \geq 1$. Thus $d_1 + \cdots + d_{k+1} = n$, and hence the number of integers in $[F_n, F_{n+1})$ with exactly $k + 1$ summands is $\binom{n+k+1-1}{k+1-1} = \binom{n+k}{k}$. This is the probability density function; in this case the proof is completed by using Stirling’s formula to see convergence to a Gaussian (the general case is harder, and proceeds via generating functions).

Returning to the gaps, our goal is to analyze the distribution of the d_j ’s. Given a decomposition

$$m = G_{i_k} + G_{i_{k-1}} + \cdots + G_{i_2} + G_{i_1}, \quad (1.2)$$

we define the gaps to be

$$i_k - i_{k-1}, \quad i_{k-1} + i_{k-2}, \quad i_2 - i_1; \quad (1.3)$$

note we do not consider $i_1 - 0$ as a gap (as there are on the order of $n(G_{n+1} - G_n)$ summands, one additional gap for each m has a negligible affect). We look at the behavior of the average gap measure. By this we mean we amalgamate all gaps from all decompositions of all integers in $[G_n, G_{n+1})$, and show that as $n \rightarrow \infty$ this average converges to a limiting distribution which decays geometrically for gap lengths larger than L , with decay ratio equal to the largest eigenvalue of the recurrence of the G_n 's. In a sequel paper [BILMT] we show that for each $m \in [G_n, G_{n+1})$ the associated gap measure of m converges almost surely to this average as $n \rightarrow \infty$ (as well as determining the limiting distribution of the longest gap in each decomposition); we concentrate here on the average gap measure for a special class of recurrence coefficients for two reasons. In addition to simplifying the proofs and highlighting the method, more can be proved about the limiting behavior in these cases, and it is thus worth isolating these results. We consider what we shall name Kangaroo recurrences.

Definition 1.1. A **Kangaroo recurrence** is a positive linear recurrence relation

$$K_{n+1} = c_1 K_n + \cdots + c_L K_L = K_n + K_{n-g} + K_{n-2g} + \cdots + K_{n-\ell g}, \quad (1.4)$$

with initial conditions $K_1 = 1$, and for $1 \leq n \leq \ell g$

$$K_{n+1} = c_1 K_n + c_2 K_{n-1} + \cdots + c_n K_1 + 1. \quad (1.5)$$

Important special cases include the Fibonacci numbers ($g = 1$ and $\ell = 1$), the Tribonacci ($g = 1$ and $\ell = 2$) and other similar recurrences. We call g the **hop** length.

The Kangaroo recurrences satisfy the conditions of [MW1, MW2] and lead to a unique legal decomposition, which is specified by the following.

Lemma 1.2. For each positive integer N there exists a unique decomposition $\sum_{i=1}^m a_i K_{m+1-i}$ with $a_1 > 0$, the other $a_i \geq 0$, and one of the following two conditions holds:

- We have $m < L$ and $a_i = c_i$ for $1 \leq i \leq m$.
- There exists $s \in \{1, \dots, L\}$ such that

$$a_1 = c_1, a_2 = c_2, \dots, a_{s-1} = c_{s-1} \text{ and } a_s < c_s, \quad (1.6)$$

$a_{s+1}, \dots, a_{s+\ell} = 0$ for some $\ell \geq 0$, and $\{b_i\}_{i=1}^{m-s-\ell}$ (with $b_i = a_{s+\ell+i}$) is either legal or empty.

For example, for the Tribonacci numbers the first terms are 1, 2, 4, 7, 13, while for $K_{n+1} = K_n + K_{n-2}$ the sequence begins 1, 2, 3, 4, 6, 9 (notice in this case we decompose 10 as $9 + 1$ and not $6 + 3 + 1$, as we can use the recurrence relation to replace $6 + 3$ with 9).

To state our main result, we first need a definition.

Definition 1.3 (Gap indicator random variables, average gaps). Consider a Kangaroo recurrence $\{K_n\}$.

- Let $G_{i,i+j}(m; n)$ equal 1 if K_i and K_{i+j} are summands in the generalized Zeckendorf decomposition of $m \in [K_n, K_{n+1})$ and K_{i+r} is not a summand for $0 < r < j$, and $G_{i,i+j}(m; n)$ is zero otherwise. Thus $G_{i,i+j}(m; n)$ is 1 if and only if $m \in [K_n, K_{n+1})$ has a gap of length j starting at K_i .
- Set

$$X_{i,i+j}(n) := \sum_{m \in [K_n, K_{n+1})} G_{i,i+j}(m; n); \quad (1.7)$$

$X_{i,i+j}(n)$ is the number of integers $m \in [K_n, K_{n+1})$ that have a gap of length j starting at K_i . Let $P_n(j)$ be the proportion of all gaps from decompositions of $m \in [K_n, K_{n+1})$ that are of length j starting at K_i .

- Let $Y(n)$ denote the total number of gaps between summands of the generalized Zeckendorf decomposition of $m \in [K_n, K_{n+1})$; thus

$$Y(n) := \sum_{i=1}^n \sum_{j=0}^n X_{i,i+j}(n); \quad (1.8)$$

note $Y(n)$ is $C_{\text{Lek}}n + O(1)$ by [MW1].

- The probability of a gap of length j , $P_n(j)$, is

$$P_n(j) := \frac{\sum_{i=1}^{n-j} X_{i,i+j}(n)}{Y(n)}. \quad (1.9)$$

Finally set

$$P(j) := \lim_{n \rightarrow \infty} P_n(j), \quad (1.10)$$

so $P(j)$ represents the limiting probability that a gap in a decomposition for an integer in $[K_n, K_{n+1})$ has length j .

Our main result is the following.

Theorem 1.4. *Let $\{K_n\}$ be a Kangaroo recurrence of ℓ hops of length g , with $\lambda_{g,\ell}$ the largest root of the characteristic polynomial, a_1 the generalized Binet constant in front of $\lambda_{g,\ell}$, and C_{Lek} the computable Lekkerkerker constant. Then*

$$P(j) = \begin{cases} 0 & \text{for } j < g \\ \left(\frac{a_1}{C_{\text{Lek}}}\right) \lambda_{g,\ell}^{-2g} \frac{\lambda_{g,\ell}^{-g} - \lambda_{g,\ell}^{-\ell g}}{1 - \lambda_{g,\ell}^{-g}} & \text{for } j = g \\ (\lambda_{g,\ell} - 1)^2 \left(\frac{a_1}{C_{\text{Lek}}}\right) \lambda_{g,\ell}^{-j} & \text{for } j > g, \end{cases} \quad (1.11)$$

which implies the probability of a gap of length j decays geometrically for $j > g$, with decay ratio equal to $\lambda_{g,\ell}$.

Remark 1.5. *As a quick consistency check, note that if $\ell = 1$ and $g = 1$ then the probability of a gap of length 1 is zero; this is of course nothing more than a restatement of the definition of a legal decomposition by the shifted Fibonacci numbers as specified by Zeckendorf and Lekkerkerker, see [Ze] and [Lek]. More is true; if $\ell = 1$ there are never gaps of length g , which must be true as if there were a gap of length g , in this case we could immediately use the recurrence relation to replace it.*

Theorem 1.4 holds in greater generality (see [BILMT]). In §2 we prove this case of the general result. As we are working with a very special recurrence, more is true, and we derive asymptotic values for $\lambda_{g,\ell}$ and hence the probabilities for ℓ and g large. Our technique is combinatorial, using inclusion-exclusion and the indicator random variables $X_{i,i+j}(m)$. As an additional example of the power of these methods, in §3 we apply these techniques to the far-difference representation of Alpert [Al] (she proved every integer can be written uniquely as a sum of signed Fibonacci summands, subject to certain constraints), determining the distribution of gaps in Theorem 3.2. We conclude with some remarks on open and related problems.

2. KANGAROO RECURRENCES

2.1. Preliminary Results. The starting point for our investigations is a generalization of Binet's formula, which gives us the growth rate of the K_n 's.

Lemma 2.1. *Let $\lambda_1, \dots, \lambda_L$ be the eigenvalues of the characteristic polynomial of a Kangaroo recurrence, ordered so that $\lambda_1 > |\lambda_2| \geq \dots \geq |\lambda_L|$. Then $\lambda_1 > 1$ and there exist constants such that*

$$K_n = a_1 \lambda_1^n + O(n^{L-2} \lambda_2^n). \quad (2.1)$$

More precisely, if $\lambda_1, \omega_2, \dots, \omega_r$ denote the distinct roots with multiplicities 1, m_2, \dots, m_r , then there are constants $a_1, a_{i,j}$ such that

$$K_n = a_1 \lambda_1^n + \sum_{i=2}^r \sum_{j=1}^{m_r} a_{i,j} n^{j-1} \omega_i^n. \quad (2.2)$$

The proof is standard. If $\lambda_1 \leq 1$ then our series is bounded, which contradicts the existence of a legal decomposition. The only difficulty is showing that there is a unique root of largest absolute value, and that this root is positive. This follows from the Perron-Frobenius Theorem and elementary algebra in the general case (or direct computation for the Fibonacci, Tribonacci and other specific recurrences); see [BILMT] for details.

Before proving Theorem 1.4 we isolate an important observation, which greatly simplifies the calculations.

Lemma 2.2. *Let $\{K_n\}$ be a Kangaroo recurrence with ℓ hops of length g . Then $P_n(j) = 0$ for all $j < g$; in other words, there cannot be a gap of length less than g .*

Proof. This follows immediately from the definition of a legal decomposition; specifically, from (1.6), which prevents us from choosing any of the next $g - 1$ summands after a chosen summand K_p . \square

Our combinatorial attack requires us to compute $X_{i,i+j}(n)$ to find $P_n(j)$. We can find $X_{i,i+j}(n)$ by counting the number of choices of the summands $\{K_1, K_2, \dots, K_n\}$ such that K_i, K_{i+j} and K_n are chosen, no summand whose index is between i and $i + j$ is chosen, and all other indices are free to be chosen subject to the requirement that we have a legal decomposition. Let $L_{i,i+j}(n)$ and $R_{i,i+j}(n)$ be the number of ways to choose a valid subset of summands from those before the gap of length j starting at K_i and after the gap (respectively). We note $L_{i,i+j}(n)$ and $R_{i,i+j}(n)$ are independent of each other when $j > g$; thus

$$X_{i,i+j}(n) = L_{i,i+j}(n) \cdot R_{i,i+j}(n). \quad (2.3)$$

As

$$K_{n+1} = K_n + K_{n-g} + \dots + K_{n-\ell g}, \quad (2.4)$$

any time we have a gap of length $j > g$ the recurrence ‘resets’ itself. Thus we have the following lemma.

Lemma 2.3. *Let $\{K_n\}$ be a Kangaroo recurrence with ℓ hops of length g . Consider all $m \in [K_n, K_{n+1})$ with a gap of length $j > g$ starting at K_i . The number of valid choices for subsets of summands before the gap, $L_{i,i+j}(n)$, is given by*

$$L_{i,i+j}(n) = K_{i+1} - K_i, \quad (2.5)$$

and the number of valid choices for subsets of summands after the gap, $R_{i,i+j}(n)$, is given by

$$R_{i,i+j}(n) = K_{n-i-j+2} - 2K_{n-i-j+1} + K_{n-i-j}. \quad (2.6)$$

Proof. To count $L_{i,i+j}(n)$, we must count the number of ways to have a legal decomposition that must have the summand K_i , and where all other summand choices beforehand are free. It is very important that $j > g$, as this means the summand at K_{i+j} does not interact with the summands earlier than K_i through the recurrence relation. Thus $L_{i,i+j}(n)$ is the same as the number of legal choices of summands from $\{K_1, K_2, \dots, K_i\}$ with K_i chosen. As each integer in $[K_i, K_{i+1})$ has a unique legal decomposition, we see $L_{i,i+j}(n)$ equals the number of elements in this interval, which is just $K_{i+1} - K_i$.

To compute $R_{i,i+j}(n)$, we need to consider how many ways we can choose summands from $\{K_{i+j}, K_{i+j+1}, \dots, K_n\}$ such that K_i and K_n are chosen and the resulting decomposition is legal; since $j > g$ the summands from K_i and earlier cannot affect our choices here. Thus our problem is equivalent to asking how many legal ways there are to choose summands from $\{K_1, K_2, \dots, K_{n-i-j+1}\}$ with $K_1, K_{n-i-j+1}$ both chosen and the rest free. There are many ways to compute this; the simplest is to note that this equals the number of legal choices where we *may or may not choose* K_1 , minus the number of legal choices where we *do not choose* K_1 . By a similar argument as above, the first count is $K_{n-i-j+2} - K_{n-i-j+1}$ (as it is the number of legal representations of a number in $[K_{n-i-j+1}, K_{n-i-j+2})$), while the second is $K_{n-i-j+1} - K_{n-i-j}$. The proof is completed by subtracting. \square

Note that in the above lemma we only compute $L_{i,i+j}(n)$ and $R_{i,i+j}(n)$ when the gap $j > g$. The reason is that if we can determine $P_n(j)$ for $j > g$, then since $P_n(j) = 0$ for $j < g$ the law of alternatives implies $P_n(g) = 1 - \sum_{j>g} P_n(j)$. Thus if we can show each limit $P(j)$ exists when $j > g$, then the limit $P(g)$ exists as well. Hence, in some sense the following lemma is not needed. We prefer to give it as it allows us to determine the ratio a_1/C_{Lek} , which we would not be able to do otherwise.

Lemma 2.4. *Let $\{K_n\}$ be a Kangaroo recurrence with ℓ hops of length g . Consider all $m \in [K_n, K_{n+1})$ with exactly b consecutive gaps of length g starting at K_i . Then $b \leq \ell - 1$ and number of valid choices for subsets of summands before the gap, $L_{i,g,b}(n)$, is given by*

$$L_{i,g,b}(n) = K_{i-g}, \quad (2.7)$$

and the number of valid choices for subsets of summands after the gap, $R_{i,g,b}(n)$, is given by

$$R_{i,g,b}(n) = K_{n-i-(b+1)g+1} - K_{n-i-(b+1)g}. \quad (2.8)$$

Proof. We must have $b \leq \ell - 1$, as if $b \geq \ell$ then we could use the recurrence relation. Remember by Lemma 2.2 the smallest gap is at least g . As we have exactly b consecutive gaps of length g starting at K_i , the first summand chosen to the *left* of K_i is at most K_{i-g-1} , while the first summand chosen to the *right* of K_{i+bg} is at least $K_{i+(b+1)g+1}$. This implies that $L_{i,g,b}(n)$ equals the number of ways to legally choose summands from $\{K_1, K_2, \dots, K_{i-g-1}\}$; however, unlike our previous case now we do not need to choose K_{i-g-1} . A straightforward calculation (or a telescoping sum) shows that this number is K_{i-g} .

Similarly, by shifting indices we find $R_{i,g,b}(n)$ is equivalent to the number of ways to legally choose summands from $\{K_{i+(b+1)g+1}, \dots, K_n\}$ with K_n chosen. Equivalently, this equals the number of legal choices of summands from $\{K_1, K_2, \dots, K_{n-i-(b+1)g}\}$ with $K_{n-i-(b+1)g}$, which is just $K_{n-i-(b+1)g+1} - K_{n-i-(b+1)g}$. \square

2.2. Proof of Theorem 1.4. We now prove our main result.

Proof of Theorem 1.4. There are three cases: $j < g$, $j = g$ and $j > g$. The first case is the easiest, as $P_n(j) = 0$ for all $j < g$ by Lemma 2.2.

We now consider $j > g$, as $j = g$ follows immediately by the law of alternatives (i.e., the probabilities must sum to 1). We need to compute $\lim_{n \rightarrow \infty} \frac{1}{Y(n)} \sum_{i=1}^{n-j} X_{i,i+j}(n)$. By Lemma 2.3,

$$X_{i,i+j}(n) = L_{i,i+j}(n) \cdot R_{i,i+j}(n) = (K_{i+1} - K_i) \cdot (K_{n-i-j+2} - 2K_{n-i-j+1} + K_{n-i-j}). \quad (2.9)$$

By Lemma 2.1,

$$K_r = a_1 \lambda_{g,\ell}^r + O(r^{L-2} \lambda_2^r), \quad (2.10)$$

with $|\lambda_2| < \lambda_{g,\ell}$. Thus

$$\begin{aligned} X_{i,i+j}(n) &= a_1 \lambda_{g,\ell}^i (\lambda_{g,\ell} - 1) (1 + o(1)) \cdot a_1 \lambda_{g,\ell}^{n-i-j} (\lambda_{g,\ell}^2 - 2\lambda_{g,\ell} + 1) (1 + o(1)) \\ &= a_1^2 (\lambda_{g,\ell} - 1)^3 \lambda_{g,\ell}^n \cdot \lambda_{g,\ell}^{-j} (1 + o(1)). \end{aligned} \quad (2.11)$$

As

$$Y(n) = (C_{\text{Lek}} n + O(1)) (K_{n+1} - K_n) = a_1 C_{\text{Lek}} n (\lambda_{g,\ell} - 1) \lambda_{g,\ell}^n (1 + o(1)) \quad (2.12)$$

and for any fixed j the sum over i is $n + O(1)$, we find

$$\begin{aligned} P_n(j) &= \frac{\sum_{i=1}^{n-j} X_{i,i+j}(n)}{Y(n)} \\ &= \frac{\sum_{i=1}^{n-j} a_1^2 (\lambda_{g,\ell} - 1)^3 \lambda_{g,\ell}^n \cdot \lambda_{g,\ell}^{-j} (1 + o(1))}{a_1 C_{\text{Lek}} n (\lambda_{g,\ell} - 1) \lambda_{g,\ell}^n (1 + o(1))} = \frac{a_1 (\lambda_{g,\ell} - 1)^2}{C_{\text{Lek}}} \lambda_{g,\ell}^{-j} (1 + o(1)), \end{aligned} \quad (2.13)$$

and the limit clearly exists for each n and each $j > g$.

We now turn to $j = g$. We avoid double counting by using the $L_{i,g,b}(n)$ and $R_{i,g,b}(n)$ from Lemma 2.4. To find the number of gaps of length g , we look at all strings of $b \leq \ell - 1$ consecutive gaps of length g over all possible starting places i , and find

$$P_n(g) = \frac{\sum_{i=1}^{n-g} \sum_{b=1}^{\ell-1} L_{i,g,b}(n) R_{i,g,b}(n)}{Y(n)}. \quad (2.14)$$

We now substitute for $L_{i,g,b}(n)$ and $R_{i,g,b}(n)$ with the values from Lemma 2.4. The proof is completed by using the generalized Binet formula (Lemma 2.1), the geometric series formula, and collecting the terms. \square

Remark 2.5. As a safety check, we confirm the theorem for the Fibonacci and Tribonacci numbers. For the Fibonacci numbers, $\lambda_1 = \phi = (1 + \sqrt{5})/2$, $a_1 = \phi/\sqrt{5}$ (remember these are the ‘shifted’ Fibonacci numbers as originated with Zeckendorf and Lekkerkerker), and $C_{\text{Lek}} = 1/(\phi^2 + 1)$. We see $P(1) = 0$ and the sum of the other probabilities is 1. For the Tribonacci numbers, $P(1)$ is no longer zero. We have $C_{\text{Lek}} = a_1(3\lambda_1^2 - 1)/(\lambda_1^3(\lambda_1^2 - 1))$, which does lead to the probabilities summing to 1.

2.3. Approximating $\lambda_{g,\ell}$ and Relevant Probability Ratios. We end this section with some quick approximations for $\lambda_{g,\ell}$ when $\ell \rightarrow \infty$.

Lemma 2.6. *For a Kangaroo recurrence of ℓ hops of length g , for large ℓ, g we have $\lambda_{g,\ell} \approx \left(1 + \frac{\alpha}{g}\right)$, where $\alpha \approx \log(g) - \log(\log(g)) + \frac{\log(\log(g))}{\log(g)}$. In particular, for large ℓ and g we have*

$$\lambda_{g,\ell} \approx 1, \quad \lambda_{g,\ell}^{-g} \approx e^{-\alpha(g)} \approx \frac{\log g}{g}. \quad (2.15)$$

Proof. We drop all lower order terms in the arguments below, as our goal is to highlight the limiting behavior. We derive a transcendental equation from the characteristic polynomial of the Kangaroo recurrence. Notice that $K_{n+1} \approx \lambda_{g,\ell} K_n$. Consider the Kangaroo recurrence $K_{n+1} = K_n + K_{n-g} + \dots + K_{n-\ell g}$. The left hand side approximately equals $\lambda_{g,\ell} K_n$, and for ℓ large the right hand side is essentially

$$K_n \sum_{m=0}^{\ell} \lambda_{g,\ell}^{-mg} \approx K_n \sum_{m=0}^{\infty} \lambda_{g,\ell}^{-mg} = \frac{K_n}{1 - \lambda_{g,\ell}^{-g}}. \quad (2.16)$$

Therefore

$$\lambda_{g,\ell} \approx \frac{1}{1 - \lambda_{g,\ell}^{-g}}, \quad (2.17)$$

which implies

$$\lambda_{g,\ell}^g - \lambda_{g,\ell}^{g-1} - 1 \approx 0. \quad (2.18)$$

From the generalized Binet formula we know that $\lambda_{g,\ell}$ grows exponentially with g . For large g we can write

$$\lambda_{g,\ell}^g = e^{\alpha(g)}, \quad \text{or} \quad \lambda_{g,\ell} = e^{\alpha(g)/g} \approx 1 + \frac{\alpha(g)}{g}. \quad (2.19)$$

Substituting into (2.18) yields

$$e^{\alpha(g)} - e^{\alpha(g)} \left(1 + \frac{\alpha(g)}{g}\right)^{-1} - 1 \approx 0, \quad (2.20)$$

which we rewrite as

$$\alpha(g) e^{\alpha(g)} \approx g. \quad (2.21)$$

By direct substitution we see that for g sufficiently large,

$$\alpha(g) \approx \log(g) - \log(\log(g)) + \frac{\log(\log(g))}{\log(g)}. \quad (2.22)$$

□

Now that we have the probability distribution for gaps for Kangaroo recurrences, we can study the ratios of the probabilities of certain events. Since gaps of length g are a threshold event, a natural question to ask is what is the probability of getting a gap of length g compared to any other gap length?

Lemma 2.7. *For large g and $\ell, n \rightarrow \infty$, the ratio of the probability of obtaining a gap of length g to a gap of length exceeding g is*

$$\frac{\text{Prob}(\text{gap of length } g)}{\text{Prob}(\text{gap of length at least } g+1)} \approx \frac{\log g}{g}. \quad (2.23)$$

Proof. From Theorem 1.4, we have

$$\frac{\text{Prob}(\text{gap of } g)}{\text{Prob}(\text{gap at least } g+1)} = \frac{\lambda_{g,\ell}^{-2g} \lambda_{g,\ell}^{-g} / (1 - \lambda_{g,\ell}^{-g})}{\lambda_{g,\ell}^{-g} (\lambda_{g,\ell} - 1)}. \quad (2.24)$$

For large g , ℓ , and n , we use Lemma 2.6 and some algebraic manipulation to deduce the claim. \square

3. FAR-DIFFERENCE REPRESENTATIONS

We now consider a natural generalization, signed Zeckendorf decompositions. Alpert [Al] proved that every positive integer has a unique representation as a sum of signed shifted Fibonacci numbers, where the gap between opposite signed summands must be at least 3 and between same signed summands must be at least 4; this is called the **far-difference** representation. Miller and Wang [MW1] proved the generalized Lekkerkerker's theorem holds here as well, and proved Gaussian behavior for the number of summands (it is a bivariate Gaussian, as there are positive and negative summands).

Our techniques generalize immediately and yield formulas for the distribution of average gaps. We concentrate on gaps between any adjacent summands, though similar reasoning would yield results restricted to gaps between same signed or opposite signed summands. Before stating and proving these generalizations, we first introduce some definitions and useful results.

Given an integer m , we write its far-difference representation by

$$m = \epsilon_{i_k} F_{i_k} + \epsilon_{i_{k-1}} F_{i_{k-1}} + \cdots + \epsilon_{i_1} F_{i_1}, \quad \epsilon_j \in \{-1, 1\}. \quad (3.1)$$

Let $\mathcal{N}(\epsilon_i F_i, \epsilon_j F_j)$ denote the number of numbers whose far-difference decomposition starts with $\epsilon_i F_i$ and ends with $\epsilon_j F_j$, where the ϵ 's are ± 1 ; similarly, let $\mathcal{N}(\epsilon_j F_j)$ be the number of numbers whose decomposition ends with $\epsilon_j F_j$. We consider integers in the interval $(S_{n-1}, S_n]$, where

$$S_n = F_n + F_{n-4} + F_{n-8} + F_{n-12} + \cdots; \quad (3.2)$$

the interval is a bit different than before as we have a signed decomposition, and have the ability to overshoot and then correct through subtraction.

Lemma 3.1. *We have*

$$\mathcal{N}(\epsilon_i F_i, \epsilon_j F_j) = \mathcal{N}(\epsilon_i F_1, \epsilon_j F_{j-i+1}) \quad (3.3)$$

and

$$\mathcal{N}(-F_1, +F_j) + \mathcal{N}(+F_1, +F_j) = \mathcal{N}(+F_j) - \mathcal{N}(+F_{j-1}). \quad (3.4)$$

Note

$$\mathcal{N}(+F_r) = S_r - S_{r-1}, \quad (3.5)$$

and by symmetry $\mathcal{N}(+F_r) = \mathcal{N}(-F_r)$.

Proof. The proof is similar to the proofs from §2, proceeding by shifting indices down for the first part, and inclusion-exclusion for the second. The final claim on $\mathcal{N}(+F_r)$ follows by noting this is the cardinality of $(S_{r-1}, S_r]$. \square

We can similarly determine the average gap behavior.

Theorem 3.2. *As $n \rightarrow \infty$, the probability $P(j)$ of a gap of length j in a far-difference decomposition of integers in $(S_{n-1}, S_n]$ converges to geometric decay for $j \geq 4$, with decay constant equal to the golden mean. Specifically, if $a_1 = \phi/\sqrt{5}$ (the Binet constant of the largest eigenvalue, $\phi = (1 + \sqrt{5})/2$), then $P(j) = 0$ if $j \leq 2$ and*

$$P(j) = \begin{cases} \frac{10a_1\phi}{\phi^4-1}\phi^{-k} & \text{if } j \geq 4 \\ \frac{5a_1}{\phi^2(\phi^4-1)} & \text{if } j = 3. \end{cases} \quad (3.6)$$

Proof. We first count gaps of length 3 in far-difference representations of integers in $(S_{n-1}, S_n]$. Let $X_{i,i+3}(n)$ be the number of representations with a gap between of length 3 from F_i to F_{i+3} (we may assume $i+3 < n-4$ and not worry about boundary effects, which are lower order). We note that since the gap length is 3, the sign of the F_i term is the opposite of the sign of the F_{i+3} term.

This gives us two cases. In the first case, we have $+F_i$ and $-F_{i+3}$ in the decomposition, while in the second case we have $-F_i$ and $+F_{i+3}$. Thus there are $\mathcal{N}(+F_i)\mathcal{N}(-F_{i+3})$ decompositions in case 1 and $\mathcal{N}(-F_i)\mathcal{N}(+F_{i+3})$ decompositions in case 2. By Lemma 3.1 the sum of the number of legal decompositions in the two cases is

$$X_{i,i+3}(n) = \mathcal{N}(+F_i) (\mathcal{N}(+F_{n-(i+2)}) - \mathcal{N}(+F_{n-(i+3)})) . \quad (3.7)$$

The proof for gaps of length 3 is completed by using $\mathcal{N}(+F_j) = S_j - S_{j-1}$, Binet's formula and the geometric series formula to show

$$S_k = \frac{a_1\phi^k}{1 - \frac{1}{\phi^4}} + O(1), \quad (3.8)$$

and the fact that the number of summands $Y(n)$ satisfies

$$Y(n) = \left(\frac{1}{5}n + \frac{366 - 118\sqrt{5}}{20} + o(1) \right) (S_n - S_{n-1}) \quad (3.9)$$

(see [MW1] for a proof of this last result). We simply substitute these results into

$$P_n(j) = \frac{1}{Y(n)} \sum_{i=1}^{n-j} X_{i,i+j}(n), \quad (3.10)$$

simplify and then take the limit as $n \rightarrow \infty$.

The case of $j \geq 4$ is easier, as now the ends' parities are independent of each other. The claim follows from a similar calculation. \square

4. CONCLUSION AND OPEN QUESTIONS

In this work we studied the average gap distribution arising from special positive linear recurrence sequences. The techniques generalize immediately to any such sequence where each recursion coefficient c_i is positive (though it is harder to get good asymptotic expansions as in §2.3); more involved formulas exist if this condition is not satisfied. It is straightforward

to see there is geometric decay for gaps larger than the recurrence length; this is essentially due to the fact that the large separation makes the left and right parts independent. Using moment techniques, in the sequel paper [BILMT] these results are extended further. They associate a gap measure to each integer in the interval $[G_n, G_{n+1})$ and show that as $n \rightarrow \infty$ almost surely each individual gap measure converges to the average gap measure. It is quite interesting that the gap problem is significantly easier than counting the number of summands; this is very different than similar problems in random matrix theory, where the eigenvalue densities of many structured ensembles are known, but not the gaps between adjacent eigenvalues (or, if known, these results are very recent and require significantly more machinery than is needed for the densities).

The combinatorial vantage here, which is an outgrowth of [KKMW, MW1], is useful for a variety of other problems. Conditioning on the number of summands is useful in investigations of the longest gap in generalized Zeckendorf decompositions [BILMT] and the asymptotic average of the number of terms in the Ostrowski α -decomposition [BCCSW]; one natural future project would be to study the distribution of gaps in the α -decomposition, as well as considering this and more general signed decompositions.

REFERENCES

- [Al] H. Alpert, *Differences of multiple Fibonacci numbers*, Integers: Electronic Journal of Combinatorial Number Theory **9** (2009), 745–749.
- [BILMT] A. Bower, R. Insoft, S. Li, S. J. Miller and P. Tosteson, *The Distribution of Gaps between Summands in Generalized Zeckendorf Decompositions*, preprint.
- [BCCSW] E. Burger, D. C. Clyde, C. H. Colbert, G. H. Shin and Z. Wang, *A Generalization of a Theorem of Lekkerkerker to Ostrowski’s Decomposition of Natural Numbers*, Acta Arith. **153** (2012), 217–249.
- [DG] M. Drmota and J. Gajdosik, *The distribution of the sum-of-digits function*, J. Théor. Nombres Bordeaux **10** (1998), no. 1, 17–32.
- [EK] P. Erdős and M. Kac, *The Gaussian Law of Errors in the Theory of Additive Number Theoretic Functions*, American Journal of Mathematics **62** (1940), no. 1/4, pages 738–742.
- [FGNPT] P. Filippini, P. J. Grabner, I. Nemes, A. Pethő, and R. F. Tichy, *Corrigendum to: “Generalized Zeckendorf expansions”*, Appl. Math. Lett., **7** (1994), no. 6, 25–26.
- [GT] P. J. Grabner and R. F. Tichy, *Contributions to digit expansions with respect to linear recurrences*, J. Number Theory **36** (1990), no. 2, 160–169.
- [GTNP] P. J. Grabner, R. F. Tichy, I. Nemes, and A. Pethő, *Generalized Zeckendorf expansions*, Appl. Math. Lett. **7** (1994), no. 2, 25–28.
- [Ho] V. E. Hoggatt, *Generalized Zeckendorf theorem*, Fibonacci Quarterly **10** (1972), no. 1 (special issue on representations), pages 89–93.
- [Ke] T. J. Keller, *Generalizations of Zeckendorf’s theorem*, Fibonacci Quarterly **10** (1972), no. 1 (special issue on representations), pages 95–102.
- [LT] M. Lamberger and J. M. Thuswaldner, *Distribution properties of digital expansions arising from linear recurrences*, Math. Slovaca **53** (2003), no. 1, 1–20.
- [Len] T. Lengyel, *A Counting Based Proof of the Generalized Zeckendorf’s Theorem*, Fibonacci Quarterly **44** (2006), no. 4, 324–325.
- [Lek] C. G. Lekkerkerker, *Voorstelling van natuurlyke getallen door een som van getallen van Fibonacci*, Simon Stevin **29** (1951-1952), 190–195.
- [KKMW] M. Koloğlu, G. Kopp, S. J. Miller and Y. Wang, *On the number of summands in Zeckendorf decompositions*, Fibonacci Quarterly **49** (2011), no. 2, 116–130.
- [MW1] S. J. Miller and Y. Wang, *From Fibonacci Numbers to Central Limit Type Theorems*, Journal of Combinatorial Theory, Series A **119** (2012), no. 7, 1398–1413.

- [MW2] S. J. Miller and Y. Wang, *Gaussian Behavior in Generalized Zeckendorf Decompositions*, to appear in the conference proceedings of the 2011 Combinatorial and Additive Number Theory Conference. <http://arxiv.org/abs/1107.2718>.
- [Ste1] W. Steiner, *Parry expansions of polynomial sequences*, *Integers* **2** (2002), Paper A14.
- [Ste2] W. Steiner, *The Joint Distribution of Greedy and Lazy Fibonacci Expansions*, *Fibonacci Quarterly* **43** (2005), 60–69.
- [Ze] E. Zeckendorf, *Représentation des nombres naturels par une somme des nombres de Fibonacci ou de nombres de Lucas*, *Bulletin de la Société Royale des Sciences de Liège* **41** (1972), pages 179–182.

E-mail address: obeckwith@gmail.com

DEPARTMENT OF MATHEMATICS, HARVEY MUDD COLLEGE, CLAREMONT, CA 91711

E-mail address: amandarg@umd.umich.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MICHIGAN-DEARBORN, DEARBORN, MI 48128

E-mail address: louis.gaudet@yale.edu

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06510

E-mail address: rinsoft@wellesley.edu

DEPARTMENT OF MATHEMATICS, WELLESLEY COLLEGE, WELLESLEY, MA 02481

E-mail address: jjl2357@berkeley.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, BERKELEY, CA 94720

E-mail address: sjm1@williams.edu, Steven.Miller.MC.96@aya.yale.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, WILLIAMS COLLEGE, WILLIAMSTOWN, MA 01267

E-mail address: Philip.D.Tosteson@williams.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, WILLIAMS COLLEGE, WILLIAMSTOWN, MA 01267